# Rhesus macaque CEACAM-5 / CD66e Protein, His Tag (MALS verified)

Catalog # CE5-R52H3



## **Synonym**

CEACAM-5,CD66e,CEA,Meconium antigen 100

#### Source

Rhesus macaque CEACAM-5 Protein, His Tag(CE5-R52H3) is expressed from human 293 cells (HEK293). It contains AA Gln 35 - Gly 685 (Accession # A0A345AMC6-1).

Predicted N-terminus: Gln 35

### **Molecular Characterization**

CEACAM-5(Gln 35 - Gly 685) A0A345AMC6-1

Poly-his

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 73.7 kDa. The protein migrates as 110-180 kDa when calibrated against <u>Star Ribbon Pre-stained Protein Marker</u> under reducing (R) condition (SDS-PAGE) due to glycosylation.

#### Endotoxin

Less than 1.0 EU per µg by the LAL method.

## **Purity**

>95% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

#### **Formulation**

Lyophilized from  $0.22~\mu m$  filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

#### Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

#### Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:


- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

## **SDS-PAGE**



Rhesus macaque CEACAM-5 Protein, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95% (With <u>Star Ribbon Pre-stained Protein Marker</u>).

## **SEC-MALS**



The purity of Rhesus macaque CEACAM-5 Protein, His Tag (Cat. No. CE5-R52H3) is more than 90% and the molecular weight of this protein is around 225-260 kDa verified by SEC-MALS.

Repor

## Background

This gene encodes a cell surface glycoprotein that represents the founding member of the carcinoembryonic antigen (CEA) family of proteins. The encoded protein is used as a clinical biomarker for gastrointestinal cancers and may promote tumor development through its role as a cell adhesion molecule. Additionally, the encoded



# Rhesus macaque CEACAM-5 / CD66e Protein, His Tag (MALS verified)

Catalog # CE5-R52H3



protein may regulate differentiation, apoptosis, and cell polarity. This gene is present in a CEA family gene cluster on chromosome 19. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2015]

**Clinical and Translational Updates** 

