



Synonym


NECTIN4, LNIR, PRR4, PVRL4

Source

Biotinylated Human Nectin-4 Protein, His,Avitag(NE4-H82E3) is expressed from human 293 cells (HEK293). It contains AA Gly 32 - Ser 243 (Accession # [Q96NY8-1](#)).

Predicted N-terminus: His

Molecular Characterization



This protein carries a polyhistidine tag at the N-terminus and an Avi tag (Avitag™) at the C-terminus.

The protein has a calculated MW of 26.7 kDa. The protein migrates as 30-35 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Labeling

Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.

Protein Ratio

Passed as determined by the HABA assay / binding ELISA.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>90% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE



Biotinylated Human Nectin-4 Protein, His,Avitag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With [Star Ribbon Pre-stained Protein Marker](#)).

Bioactivity-ELISA

SEC-MALS




The purity of Biotinylated Human Nectin-4 Protein, His,Avitag (Cat. No. NE4-H82E3) is more than 85% and the molecular weight of this protein is around 25-35 kDa verified by SEC-MALS.

Report

Discounts, Gifts,
and more!





Immobilized Anti-Nectin-4 Antibody, Mouse IgG1 at 2 µg/mL (100 µL/well) can bind Biotinylated Human Nectin-4 Protein, His,Avitag (Cat. No. NE4-H82E3) with a linear range of 0.1-2 ng/mL (QC tested).

Background

Nectin-4(Nectin cell adhesion molecule 4) is a Nectin and nectin-like family of cell adhesion molecules. It is found in high levels in normal embryonic and fetal tissues, but decreases after adulthood and is limited in distribution in healthy tissues. Nectin-4 is overexpressed in a variety of tumor cells and promotes tumor cell growth and proliferation. Solid tumors with high expression of Nectin-4 include bladder cancer, pancreatic cancer, triple negative breast cancer, non-small cell lung cancer, stomach cancer, esophageal cancer, ovarian cancer and many other cancers. Studies have found that Nectin-4 promotes tumor cell proliferation, differentiation, migration and invasion by activating the PI3K/Akt pathway, and is believed to play an important role in the occurrence and metastasis of cancer. It is based on these research backgrounds that drug research targeting this target has emerged.

Clinical and Translational Updates

Discounts, Gifts,
and more!

